

# Prevention of Infection in Open Fractures

Charalampos G. Zalavras, MD, PhD, FACS

## KEYWORDS

- Open fracture • Infection • Antibiotics • Debridement • Soft tissue coverage
- Fracture fixation

## KEY POINTS

- Open fractures are challenging injuries and are associated with increased risk for complications, such as infection.
- The goals of open fracture management are prevention of infection, fracture union, and restoration of function.
- Management principles include careful patient and injury assessment, early administration of antibiotics, thorough surgical debridement, wound management with soft tissue coverage, and stable fracture fixation.

## INTRODUCTION

Open fractures are often the result of high-energy trauma and may be associated with life-threatening injuries.<sup>1</sup> Open fractures are characterized by definition by soft tissue injury that results in communication of the fracture site with the outside environment and contamination of the fracture site with microorganisms or even introduction of foreign bodies into the wound.<sup>2</sup> Moreover, depending on the severity of injury, there is damage to the surrounding soft tissue envelope and to bone vascularity, which compromises the healing potential, as well as the response of the host defense mechanisms to contaminating microorganisms. As a result, open fractures are associated with increased risk for complications, such as infection and nonunion, and present a challenging problem to the treating physician.

The goals of open fracture management are prevention of infection, fracture union, and restoration of function. These goals are best achieved by careful patient and injury evaluation, early administration of systemic antibiotics supplemented by local delivery of antibiotics in severe injuries, thorough surgical debridement, wound management

---

Disclosure Statement: The author has nothing to disclose.

LAC+USC Medical Center, Keck School of Medicine, University of Southern California, 1200 North State Street, GNH 3900, Los Angeles, CA 90033, USA

E-mail address: [zalavras@usc.edu](mailto:zalavras@usc.edu)

Infect Dis Clin N Am ■ (2017) ■–■

<http://dx.doi.org/10.1016/j.idc.2017.01.005>

0891-5520/17/© 2017 Elsevier Inc. All rights reserved.

[id.theclinics.com](http://id.theclinics.com)

with soft tissue coverage if needed, and stable fracture fixation. Key principles in prevention of infection following open fractures are summarized in **Box 1**.

## EVALUATION AND CLASSIFICATION

### *Evaluation of Patient and Extremity*

Open fractures are often caused by high-energy trauma, such as motor vehicle, motorcycle, and auto versus pedestrian injuries, and may be associated with potentially life-threatening injuries to the head, chest, and abdomen.<sup>1,3</sup> The injury severity score of patients with open diaphyseal fractures of the femur is 18.1 on average.<sup>1</sup> Thorough evaluation of every patient with an open fracture is necessary to diagnose and treat associated injuries, and appropriate resuscitation measures should be initiated on presentation of the patient.

The neurovascular status of the injured extremity should be carefully assessed and documented. The size, location, and degree of gross contamination of the open fracture wound should be evaluated. The wound should be irrigated, gross contamination should be removed, and a sterile dressing should be applied. The treating surgeon should not forget that the complication of compartment syndrome may still develop, despite the presence of the open fracture wound, especially in injuries with a severe crushing component.<sup>4</sup> The fractured extremity should be grossly realigned and immobilized with a splint. Intravenous antibiotic therapy should be started and tetanus prophylaxis should be given depending on the patient's immunization status. Fracture characteristics, such as location, articular involvement, and comminution, should be assessed by imaging studies to plan fixation of the fracture.

### *Classification of Open Fractures*

Open fractures vary in severity depending on the mechanism and energy of injury. Therefore, classification systems of open fractures have been developed to describe the injury, guide treatment, determine prognosis, and compare various treatment methods for research purposes. The classification system of Gustilo and Anderson, subsequently modified by Gustilo, Mendoza, and Williams, has been extensively used (**Box 2**).<sup>5,6</sup> Newer classification systems have also been proposed.<sup>7</sup>

#### **Box 1**

##### **Key principles in prevention of infection after open fractures**

- Detailed evaluation of patient for associated and potentially life-threatening injuries.
- Systemic antibiotic therapy on patient presentation.
- Intraoperative assessment of severity of injury and classification of open fracture.
- Thorough surgical debridement with removal of all devitalized tissue and foreign bodies.
- Local antibiotic delivery with the bead-pouch technique in severe injuries.
- Primary wound closure is an option for less-severe injuries if only healthy, viable tissue is present in the wound after a meticulous debridement.
- Delayed closure with second-look debridement for more severe injuries.
- Local or free muscle flaps for extensive soft tissue damage.
- Fracture stabilization with appropriate technique based on fracture, soft tissue, and patient characteristics.

**Box 2****Gustilo and Anderson classification of open fractures**

- Type I: Wound of 1 cm or smaller, with minimal contamination or muscle crushing.
- Type II: Wound more than 1 cm long with moderate soft tissue damage and crushing. Bone coverage is adequate and comminution is minimal.
- Type IIIA: Extensive soft tissue damage, often due to a high-energy injury with a crushing component. Massively contaminated wounds and severely comminuted or segmental fractures are included in this subtype. Bone coverage is adequate.
- Type IIIB: Extensive soft tissue damage with periosteal stripping and bone exposure, usually with severe contamination and bone comminution. Flap coverage is required.
- Type IIIC: Arterial injury requiring repair.

The severity of the open fracture determines the risk of infection, which ranges from 0% to 2% for type I open fractures, 2% to 10% for Type II, and 10% to 50% for type III fractures.<sup>6,8</sup>

As with all classification systems, reliability of the classification and agreement among observers remains an issue. A study evaluating the responses of orthopedic surgeons who were asked to classify open fractures of the tibia on the basis of videotaped case presentations reported that the average agreement among observers was 60%.<sup>9</sup>

It is important to remember that the true extent and severity of the injury cannot be accurately assessed in the emergency department. The degree of contamination and soft tissue crushing are important factors for classifying an open fracture that may be mistakenly overlooked in a wound of small size. Therefore, classification of the fracture should be done in the operating room following wound exploration and debridement.

## ANTIBIOTIC ADMINISTRATION

Early antibiotic administration is a key principle of open fracture management, because most patients with open fractures have wounds contaminated with microorganisms.<sup>5,10</sup> Systemic antibiotics should be administered in all patients with open fractures and local antibiotics in select cases.

### *Role of Antibiotics in Reducing Infection*

The role of antibiotics in reducing the infection rate in patients with open fractures was demonstrated by Patzakis and colleagues<sup>10</sup> in a prospective randomized study. The infection rate when cephalothin was administered before debridement was 2.3% (2 of 84 fractures) compared with 13.9% (11 of 79 fractures) when no antibiotics were used.<sup>10</sup>

### *Timing of Antibiotics*

Early antibiotic administration is very important. Patzakis and Wilkins<sup>8</sup> initially reported in 1989 that a delay longer than 3 hours from injury was associated with increased risk of infection. Subsequent animal and clinical studies have corroborated these findings. Penn-Barwell and colleagues<sup>11</sup> used a rat femur model with a defect contaminated with *Staphylococcus aureus* and found that delaying antibiotics to 6 or 24 hours had a detrimental effect on the infection rate regardless of the timing of surgery. Lack and colleagues<sup>12</sup> in a retrospective study of 137 type III open tibia factors

demonstrated that administration of antibiotics beyond 66 minutes from injury was an independent risk factor for infection with an odds ratio of almost 4.

### **Choice of Antibiotics**

---

Although the necessity of antibiotics has been definitively established, there is no consensus on the optimal choice of agents and specifically on the necessity of gram-negative coverage, especially for less-severe injuries.

In severe (type III) open fractures there is wide agreement that gram-positive and gram-negative coverage is needed and this is usually provided by a first-generation cephalosporin, such as cefazolin, and by an aminoglycoside, such as gentamicin.<sup>8,13–16</sup> However, others have advocated against aminoglycoside use and gram-negative coverage.<sup>17</sup>

In less-severe (type I and II) open fractures some investigators have recommended coverage only for gram-positive organisms with administration of a cephalosporin.<sup>13–15</sup> In contrast, combined gram-positive and gram-negative coverage for these less-severe open fractures has been proposed by other investigators to provide coverage against contaminating gram-negative organisms.<sup>8,16</sup>

Patzakis and Wilkins<sup>8</sup> reported that combination therapy in open tibia fractures reduced the infection rate (4.5%, 5 of 109) compared with cephalosporin only (13%, 25 of 192). This study did not analyze type I and II open fractures separately but the distribution of fracture types was comparable between the 2 groups that received combination therapy versus cephalosporin only.

It is important to remember that the severity of injury will be better appreciated in the operating room following wound exploration and debridement and an open fracture may be misclassified in the emergency department. As a result, a type IIIA open fracture with a wound of small size may be misclassified as type I or II open fracture and treated with cephalosporin only. Interestingly, a study of 189 patients with 202 open fractures, most of which were treated without gram-negative coverage, reported that gram-negative organisms were isolated in 55% of the infections that developed during follow-up.<sup>18</sup>

The potential nephrotoxicity of aminoglycosides is a concern that may explain the reluctance to administer aminoglycosides in the setting of open fractures. A recent study evaluated patients with open fractures who were treated either with cefazolin ( $n = 41$ ) or with cefazolin and gentamicin ( $n = 113$ ).<sup>19</sup> Baseline characteristics and risk factors for renal dysfunction did not vary between groups. There was no difference in the development of acute kidney injury (4.8% in patients receiving cefazolin vs 4% patients receiving cefazolin and gentamicin).<sup>19</sup> This may be, in part, due to the short duration of aminoglycosides used.

Quinolones have been proposed as an oral alternative to intravenous antibiotics, based on their broad-spectrum antimicrobial coverage, bactericidal properties, oral administration, and good tolerance. A randomized prospective study showed that ciprofloxacin as a single agent in type I and II open fractures resulted in a similar infection rate (6%) compared with combination therapy with cefamandole and gentamicin.<sup>20</sup> However, in type III open fractures, ciprofloxacin was associated with a higher infection rate of 31% compared with 7.7% in the combination therapy group. Moreover, quinolones have been associated with inhibition of osteoblasts.<sup>21</sup> Quinolones are excellent agents against gram-negative pathogens and are considered an alternative when aminoglycosides cannot be used.

A recent study compared use of piperacillin/tazobactam versus cefazolin plus gentamicin for antibiotic prophylaxis in patients with type 3 open fractures and found similar rates of surgical site infections at 1 year (31.4% vs 32.4%, respectively).<sup>22</sup>

Piperacillin/tazobactam has a spectrum of activity similar to that of a cephalosporin plus an aminoglycoside, but its advantages include administration of a single agent and coverage against anaerobes in cases of fecal or clostridial contamination that may occur with farm-related injuries.

Anaerobic coverage (eg, penicillin, clindamycin, or metronidazole) should be added in injuries with potential contamination with clostridial organisms (eg, farm injuries) and in vascular injuries that can create conditions of ischemia and low oxygen tension. The goal is to prevent clostridial myonecrosis (gas gangrene). However, prevention of this catastrophic and life-threatening complication is primarily based on thorough surgical debridement. Antibiotic therapy is not a substitute for inadequate debridement.

Antimicrobial resistance in bacteria has been emerging and infections due to methicillin-resistant *S aureus* (MRSA), multidrug-resistant gram-negative bacilli, and vancomycin-resistant *Enterococci* are progressively increasing. This creates concerns about the adequacy of current antibiotic protocols, especially against MRSA. On the other hand, expanded prophylactic coverage with vancomycin may facilitate the emergence of glycopeptide-resistant organisms. A randomized controlled trial compared administration of a combination of vancomycin and cefazolin to administration of only cefazolin in 101 patients with open fractures and found no difference in the infection rates between the group receiving vancomycin and cefazolin (19%) versus the group receiving only cefazolin (15%).<sup>23</sup> As a result, the routine use of vancomycin in open fractures cannot be recommended based on available data. Antibiotic choices are summarized in [Table 1](#).

### **The Role of Wound Cultures**

Wound cultures obtained at patient presentation or intraoperatively do not help select the optimal antibiotic regimen, because they fail to identify the organism causing a subsequent infection in most cases.<sup>24,25</sup> Only 18% (3 of 17) of infections that developed in a series of 171 open fractures were caused by an organism identified by perioperative cultures.<sup>20</sup> Another study reported that predebridement cultures identified the infecting organism only 22% of the time and postdebridement cultures 42% of

**Table 1**  
Choice of antibiotics in open fractures

Severity of Open Fracture	Proposed Coverage	Antibiotic Options
Type I, II	<ul style="list-style-type: none"> <li>• Gram-positive organisms (some investigators)</li> <li>OR</li> <li>• Gram-positive and gram-negative organisms (some investigators)</li> </ul>	<ul style="list-style-type: none"> <li>• Cephalosporin (1st generation)</li> <li>OR</li> <li>• Cephalosporin (1st generation) and aminoglycoside (quinolone as alternative for gram-negative coverage)</li> </ul>
Type III	<ul style="list-style-type: none"> <li>• Gram-positive and gram-negative organisms (some investigators)</li> </ul>	<ul style="list-style-type: none"> <li>• Cephalosporin (1st generation) and aminoglycoside (quinolone as alternative for gram-negative coverage)</li> <li>OR</li> <li>• Piperacillin/tazobactam</li> </ul>
Potential contamination with clostridial organisms OR vascular injuries	<ul style="list-style-type: none"> <li>• Additional coverage for anaerobic organisms</li> </ul>	<ul style="list-style-type: none"> <li>• Penicillin OR clindamycin OR metronidazole</li> </ul>

the time.<sup>25</sup> Postdebridement cultures would be preferable to predebridement cultures to select the best agents in the case of an early infection, but their accuracy is still sub-optimal. In most cases, infections are not caused by the organisms initially present in the wound but by nosocomial organisms that subsequently contaminate the wound.

### ***Duration of Systemic Antibiotic Therapy***

---

The optimal length of antibiotic therapy remains controversial. The recommended duration of antibiotic therapy is 3 days,<sup>8,15</sup> although a study comparing 1 day to 5 days of antibiotics reported similar infection rates and suggested that 1 day of antibiotics may be an option.<sup>26</sup> An additional 3-day administration of antibiotics is recommended for subsequent surgical procedures, such as repeat debridement and wound coverage.<sup>8,15</sup>

### ***Local Antibiotic Therapy***

---

Local delivery of antibiotic therapy used as an adjunct to systemic antibiotic therapy in the treatment of open fractures.<sup>27</sup> A commonly used delivery vehicle is polymethylmethacrylate (PMMA) cement, which can be manually molded by the surgeon to bead-resembling spheres with a diameter ranging from 5 to 10 mm or to spacer blocks of larger size. Bioabsorbable delivery vehicles, such as calcium sulfate, appear to be a promising alternative in clinical studies.<sup>28</sup> In a rat model of a contaminated bone defect, a bioabsorbable phospholipid gel resulted in lower infection rate compared with PMMA beads.<sup>29</sup>

An antibiotic appropriate for local delivery must be available in powder form because antibiotics in aqueous solution inhibit the polymerization process, must be heat stable to withstand the temperatures generated during the exothermic polymerization reaction, and should be active against the targeted microorganisms. Several antibiotics, including aminoglycosides, vancomycin, and cephalosporins, can be successfully incorporated into PMMA cement for local delivery. In open fractures, aminoglycosides are commonly used due to their broad spectrum of activity, heat stability, and low allergenicity.

The process of release of antibiotic from the delivery vehicle to the surrounding tissues is called elution. The difference in concentration of antibiotics between the delivery system and its environment is the key factor determining elution. Elution is facilitated by an increased surface-area-to-volume ratio of the delivery vehicle,<sup>30</sup> and by a high concentration of antibiotic in the beads or spacer.<sup>31</sup> The type of antibiotic plays a role and tobramycin has superior elution properties when compared with vancomycin.<sup>32</sup> A fluid medium is necessary for elution of antibiotics and the rate of fluid turnover influences the local antibiotic concentration. Elution of antibiotics from PMMA beads follows a biphasic pattern, with an initial rapid phase and a secondary slow phase.<sup>33</sup>

Antibiotic-impregnated PMMA beads are commercially available in Europe, but no similar product has been approved by the Food and Drug Administration for use in the United States. The delivery system has to be prepared in the operating room by the surgeon immediately before use. The PMMA powder is mixed with the antibiotic in powder form, is polymerized, and then formed into beads, which are incorporated on a 24-gauge wire. Usually 3.6 g of tobramycin are mixed with 40 g of PMMA cement. Antibiotic-impregnated PMMA beads are inserted in the open fracture wound, which is subsequently sealed by a semipermeable barrier so that the eluted antibiotic remains at the involved area to achieve a high local concentration.

The bead-pouch technique has unique advantages. First, it maintains antibiotics within the wound and achieves a high local concentration of antibiotics.<sup>34</sup> Second,

the systemic concentration remains low, thus minimizing the adverse effects of aminoglycosides. The sealing of the wound from the external environment by the semipermeable barrier prevents secondary contamination by nosocomial pathogens and may safely extend the period for soft tissue transfers. At the same time, the semipermeable barrier establishes an aerobic wound environment, which is important for avoiding catastrophic anaerobic infections. Finally, it promotes patient comfort by avoiding painful dressing changes.

The antibiotic bead-pouch technique has been shown to reduce the infection rate when used in addition to systemic antibiotics for management of open fractures. Ostermann and colleagues<sup>35</sup> compared systemic antibiotics alone with combined treatment with both systemic antibiotics and the bead-pouch technique in a series of 1085 open fractures. The infection rate was considerably lower in the antibiotic bead-pouch group (3.7% [31 of 845 fractures] vs 12% [29 of 240 fractures]) but the reduction of infection was statistically significant in type III fractures only (6.5% for the antibiotic bead pouch vs 20.6% for systemic antibiotics alone).<sup>35</sup> A systematic review reported that the absolute rate of infection was lower for all types of open tibia fractures treated with intramedullary nailing when local antibiotics were administered in addition to systemic antibiotics.<sup>36</sup>

## DEBRIDEMENT AND IRRIGATION

### *Debridement*

Thorough surgical debridement plays a critical role in the management of open fractures.<sup>37,38</sup> Devitalized tissue and foreign material promote the growth of microorganisms, constitute a barrier for the host's defense mechanisms, and should be removed. Debridement should be performed in the operating room. Surgical extension of the wound allows assessment of the degree of soft tissue damage and contamination. Skin and subcutaneous tissue are sharply debrided back to bleeding edges. Muscle is debrided until bleeding is visualized. Viable muscle can be identified by its bleeding, color, and contractility. Bone fragments without soft tissue attachments are avascular and should be discarded. Articular fragments, however, should be preserved even if they have no attached blood supply, provided they are large enough and the surgeon believes reconstruction of the involved joint is possible. A repeat debridement can be performed after 24 to 48 hours based on the degree of contamination and soft tissue damage. The goal is a clean wound with viable tissues and no infection. In injuries requiring flap coverage, debridement should be also repeated at the time of the soft tissue procedure.

### *Timing of Debridement*

Traditional concerns that delays in surgical management beyond 6 hours would lead to increased infection rates have not been substantiated in the literature.<sup>8,39–43</sup> Patzakis and Wilkins<sup>8</sup> reported in 1989 that the infection rate was similar in open fracture wounds debrided within 12 hours from injury (6.8%, 27 of 396) and in those debrided after 12 hours from injury (7.1%, 50 of 708) and concluded that elapsed time from injury to debridement is not a critical factor for development of infection in patients receiving antibiotic therapy.

Pollak and colleagues<sup>40</sup> and the LEAP study group found no relationship between time to surgical debridement and infection in 307 patients with severe open lower extremity fractures. The infection rate was 28%, 29%, and 26% in patients who underwent debridement earlier than 5 hours, 5 to 10 hours, and more than 10 hours from injury, respectively. Weber and colleagues,<sup>43</sup> in a prospective cohort study of 736

patients with open fractures, found no association between development of deep infection and time to debridement. In multivariate analysis, the only factors associated with infection were the severity (type III vs type I) and anatomic location (tibia vs upper extremity) of the fracture.

Although bacterial populations in an untreated contaminated wound increase over time, it appears that early antibiotic administration and thorough surgical debridement can effectively reduce the contamination present. As a result, small delays in surgical management do not appear to translate in increased infection rates and may allow for stabilization and resuscitation of the patient, as well as for treatment of the patient by experienced surgical teams with all necessary equipment available.

### ***Irrigation***

---

Irrigation mechanically removes foreign bodies and reduces the bacterial concentration.

Antiseptic solutions may be toxic to host cells and are better avoided.<sup>44</sup> A study comparing bacitracin solution to nonsterile castile soap solution for irrigation of open fractures found no difference in infection and nonunion rates, but an increased rate of wound-healing problems with bacitracin.<sup>45</sup> Irrigation with normal saline was not evaluated in this study.

High-pressure pulsatile lavage is more effective than low-pressure lavage in removing adherent bacteria if more than 6 hours have elapsed since the injury,<sup>46</sup> but it has been associated in experimental studies with bacterial seeding into the intramedullary canal,<sup>46</sup> with increased wound bacterial counts at 48 hours after irrigation,<sup>47</sup> and with adverse effects on early new bone formation.<sup>48</sup>

A recent randomized controlled trial of 2447 patients with open fractures compared the effects of different irrigation solutions (castile soap vs normal saline) and irrigation pressures (high pressure vs low pressure vs very low pressure) on the reoperation rate within 12 months from after the index surgery.<sup>49</sup> Irrigation with saline was associated with a significantly higher reoperation rate compared with castile soap (14.8% vs 11.6%). The reoperation rates were similar regardless of irrigation pressure, indicating that very low pressure (gravity) is an acceptable low-cost alternative method of irrigation.<sup>49</sup>

The author of the current article recommends irrigating the wound with 9 L of saline by gravity tubing.

## **WOUND MANAGEMENT**

### ***Wound Closure***

---

The optimal time for wound closure remains controversial.<sup>50</sup> Primary wound closure generates concerns for potential development of the catastrophic complication of gas gangrene (clostridial myonecrosis) that may lead to loss of the limb or even death of the patient.<sup>51</sup> However, gas gangrene has mostly complicated military wounds with severe tissue injury, gross contamination, and inadequate antibiotic therapy or debridement.

Clinical studies of civilian injuries have not shown an increased infection rate following primary closure,<sup>8,52–54</sup> and suggested that primary closure may prevent secondary contamination and reduce surgical morbidity, hospital stay, and cost.<sup>52,53</sup> Patzakis and Wilkins<sup>8</sup> reported in 1989 that primary closure did not result in increased infection rate. Specifically, infection complicated 10.6% of wounds closed primarily compared with 13.4% of wounds closed with a delay.<sup>8</sup>

Primary closure of open fracture wounds is a viable option in carefully selected cases, provided there is no severe tissue damage and contamination, especially



with soil or fecal matter, early administration of antibiotics has taken place, a meticulous debridement has been executed by an experienced surgeon, and the wound edges can be approximated without tension.

Partial wound closure is another option for less-severe injuries.<sup>55</sup> The surgical extension of the wound, created to assess the bone and soft tissues and facilitate debridement, can be closed primarily in type I and II open fractures, leaving only the injury wound open, to be closed in delayed fashion.

Delayed wound closure prevents anaerobic conditions in the wound, permits drainage, allows for repeat debridement at 24 to 48 hours, gives time to tissues of questionable viability to declare themselves, and facilitates use of the antibiotic bead-pouch technique. Delayed wound closure is recommended in injuries with extensive soft tissue damage and gross contamination, in patients presenting with a considerable delay, in wounds with tissues of questionable viability at the end of the debridement, and in wounds that cannot be approximated without tension.

If there is any doubt about the viability of the tissues and/or the adequacy of the debridement, the wound should not be closed primarily and a second-look debridement should be undertaken with the plan to perform delayed wound closure or a soft tissue coverage procedure, depending on the status of the soft tissue envelope.

In cases of delayed wound coverage, the wound should not be left exposed to the outside environment to prevent contamination with nosocomial pathogens. Instead, the antibiotic bead-pouch technique,<sup>35</sup> or negative-pressure wound therapy,<sup>56</sup> should be used.

### ***Soft Tissue Reconstruction***

---

When extensive damage to the soft tissues is present, as in type IIIB open fractures, adequate coverage may not be possible and soft tissue reconstruction is required. A well-vascularized soft tissue envelope promotes fracture healing, delivery of antibiotics, and action of the host defense mechanisms. It provides durable coverage, which prevents secondary contamination of the wound and desiccation of exposed anatomic structures, such as bone, cartilage, and tendons. The selection of coverage depends on the location and magnitude of the soft tissue defect.<sup>57,58</sup>

Soft tissue reconstruction is achieved with local or free tissue transfers. A microvascular surgeon should evaluate an open fracture with extensive soft tissue damage and participate in its management. Local pedicle muscle flaps include the gastrocnemius for proximal third tibia fractures and the soleus for middle third fractures. In distal third tibia fractures, free muscle flaps, such as the latissimus dorsi, the rectus abdominis, and the gracilis muscle, are necessary.<sup>57,58</sup> Free muscle flaps may be considered even in more proximally located fractures. Local muscles usually participate in the zone of injury; thus, their viability may be compromised. Pollak and colleagues<sup>58</sup> concluded that use of a free flap in limbs with a severe osseous injury was associated with fewer wound complications necessitating operative treatment compared with a rotational flap.

Soft tissue reconstruction should be performed early, within the first week from injury, because delays have been associated with increased flap and infectious complications.<sup>12,24,59,60</sup> Godina<sup>60</sup> advocated flap coverage within 72 hours. Gopal and colleagues<sup>57</sup> also used an early aggressive protocol in type IIIB and IIIC open fractures and observed deep infection in 6% (4/63) of fractures that were covered within 72 hours, compared with 29% (6/21) in fractures covered beyond 72 hours. It should be noted that in these studies the antibiotic bead pouch was not used, and therefore secondary contamination may have been an important confounding factor contributing to infectious complications.

## FRACTURE FIXATION

Stabilization of the open fracture is an important part of management. Stability at the fracture site prevents further injury to the soft tissues, and enhances the host response to contaminating organisms.<sup>61</sup> In addition, fracture stability facilitates wound and patient care, and allows early motion and functional rehabilitation of the extremity. Fracture stabilization can be definitive or provisional and can be accomplished with intramedullary nailing, plate and screw fixation, or external fixation. Selection among these options depends on careful evaluation of fracture, soft tissue, and patient characteristics.<sup>16</sup> More than one method may be applicable to a specific injury and the surgeon's expertise and availability of implants should also be taken into account.

Intramedullary nailing is an effective method of stabilization of diaphyseal fractures of the lower extremity.<sup>62,63</sup> Statically interlocked intramedullary nailing maintains length and alignment of the fracture bone, is biomechanically superior to other methods, and does not interfere with soft tissue management. However, cortical circulation is disrupted to a variable degree, especially when reaming of the medullary canal takes place.<sup>64</sup> An animal study showed that, despite the adverse effect of reaming on endosteal perfusion, perfusion of callus and early strength of union were similar following intramedullary nailing with or without reaming.<sup>65</sup>

Plate and screw fixation is useful in intra-articular and metaphyseal fractures because it allows anatomic reduction and restoration of joint congruency.<sup>66,67</sup> Plate and screw fixation is recommended for open diaphyseal fractures of the forearm and humerus unless there is severe muscle damage and contamination.<sup>68</sup>

External fixation can be applied in a technically easy, safe, and expedient way, with minimal blood loss. For this reason, it can be beneficial in damage-control situations, such as type IIIC open fractures and unstable polytrauma patients.<sup>69</sup> External fixation preserves the vascularity of the fracture site and avoids implant insertion at the zone of injury. Therefore, it may be useful in wounds with severe soft tissue damage and gross contamination, as in type IIIB open fractures.<sup>70,71</sup> External fixation also can be useful as provisional joint-spanning fixation in open periarticular fractures followed by definitive fixation at a second stage.<sup>72</sup>

## SUMMARY

Open fractures are challenging, often high-energy injuries that require a principle-based approach that starts with detailed evaluation of the patient for associated and potentially life-threatening injuries. The severity of the open fracture is best assessed intraoperatively and is an important factor in determining the rate of complications, such as infection and nonunion.

Systemic antibiotic therapy should be initiated on patient presentation. Local antibiotic delivery with the bead-pouch technique is beneficial in severe injuries.

Thorough surgical debridement with removal of all devitalized tissue and foreign bodies is critical for prevention of infection.

Primary wound closure is an option for less-severe injuries if only healthy, viable tissue is present in the wound after a meticulous debridement performed by an experienced surgeon. Delayed closure with a second-look debridement is recommended for more severe injuries. In the presence of extensive soft tissue damage, local or free muscle flaps should be transferred to achieve coverage. The open fracture should be stabilized using one of the available options based on fracture, soft tissue, and patient characteristics, and on surgeon's expertise. A management plan guided by the previously discussed principles will help prevent infection in these challenging injuries.

## REFERENCES

1. Court-Brown CM, Bugler KE, Clement ND, et al. The epidemiology of open fractures in adults. A 15-year review. *Injury* 2012;43(6):891–7.
2. Zalavras CG, Patzakis MJ. Open fractures: evaluation and management. *J Am Acad Orthop Surg* 2003;11(3):212–9.
3. Gustilo RB. Management of open fractures. An analysis of 673 cases. *Minn Med* 1971;54(3):185–9.
4. Blick SS, Brumback RJ, Poka A, et al. Compartment syndrome in open tibial fractures. *J Bone Joint Surg Am* 1986;68(9):1348–53.
5. Gustilo RB, Anderson JT. Prevention of infection in the treatment of one thousand and twenty-five open fractures of long bones: retrospective and prospective analyses. *J Bone Joint Surg Am* 1976;58(4):453–8.
6. Gustilo RB, Mendoza RM, Williams DN. Problems in the management of type III (severe) open fractures: a new classification of type III open fractures. *J Trauma* 1984;24(8):742–6.
7. Evans AR, Agel J, Desilva GL. A new classification scheme for open fractures. *J Orthop Trauma* 2010;24(8):457–64.
8. Patzakis MJ, Wilkins J. Factors influencing infection rate in open fracture wounds. *Clin Orthop Relat Res* 1989;(243):36–40.
9. Brumback RJ, Jones AL. Interobserver agreement in the classification of open fractures of the tibia. The results of a survey of two hundred and forty-five orthopaedic surgeons. *J Bone Joint Surg Am* 1994;76(8):1162–6.
10. Patzakis MJ, Harvey JP Jr, Ivler D. The role of antibiotics in the management of open fractures. *J Bone Joint Surg Am* 1974;56(3):532–41.
11. Penn-Barwell JG, Murray CK, Wenke JC. Early antibiotics and debridement independently reduce infection in an open fracture model. *J Bone Joint Surg Br* 2012;94(1):107–12.
12. Lack WD, Karunakar MA, Angerame MR, et al. Type III open tibia fractures: immediate antibiotic prophylaxis minimizes infection. *J Orthop Trauma* 2015;29(1):1–6.
13. Hoff WS, Bonadies JA, Cachecho R, et al. East practice management guidelines work group: update to practice management guidelines for prophylactic antibiotic use in open fractures. *J Trauma* 2011;70(3):751–4.
14. Luchette FA, Bone LB, Born CT, et al. Practice management guidelines for prophylactic antibiotic use in open fractures. 2000. Available at. <http://www.east.org/education/practice-management-guidelines/open-fractures-prophylactic-antibiotic-use-in-update>.
15. Templeman DC, Gulli B, Tsukayama DT, et al. Update on the management of open fractures of the tibial shaft. *Clin Orthop Relat Res* 1998;(350):18–25.
16. Zalavras CG, Marcus RE, Levin LS, et al. Management of open fractures and subsequent complications. *J Bone Joint Surg Am* 2007;89(4):884–95.
17. Hauser CJ, Adams CA Jr, Eachempati SR, et al. Surgical Infection Society guideline: prophylactic antibiotic use in open fractures: an evidence-based guideline. *Surg Infect (Larchmt)* 2006;7(4):379–405.
18. Chen AF, Schreiber VM, Washington W, et al. What is the rate of methicillin-resistant *Staphylococcus aureus* and Gram-negative infections in open fractures? *Clin Orthop Relat Res* 2013;471(10):3135–40.
19. Pannell WC, Banks K, Hahn J, et al. Antibiotic related acute kidney injury in patients treated for open fractures. *Injury* 2016;47(3):653–7.

20. Patzakis MJ, Bains RS, Lee J, et al. Prospective, randomized, double-blind study comparing single-agent antibiotic therapy, ciprofloxacin, to combination antibiotic therapy in open fracture wounds. *J Orthop Trauma* 2000;14(8):529–33.
21. Holtom PD, Pavkovic SA, Bravos PD, et al. Inhibitory effects of the quinolone antibiotics trovafloxacin, ciprofloxacin, and levofloxacin on osteoblastic cells in vitro. *J Orthop Res* 2000;18:721–7.
22. Redfern J, Wasilko SM, Groth ME, et al. Surgical site infections in patients with type 3 open fractures: comparing antibiotic prophylaxis with cefazolin plus gentamicin versus piperacillin/tazobactam. *J Orthop Trauma* 2016;30(8):415–9.
23. Saveli CC, Morgan SJ, Belknap RW, et al. Prophylactic antibiotics in open fractures: a pilot randomized clinical safety study. *J Orthop Trauma* 2013;27(10):552–7.
24. Fischer MD, Gustilo RB, Varecka TF. The timing of flap coverage, bone-grafting, and intramedullary nailing in patients who have a fracture of the tibial shaft with extensive soft-tissue injury. *J Bone Joint Surg Am* 1991;73(9):1316–22.
25. Lee J. Efficacy of cultures in the management of open fractures. *Clin Orthop Relat Res* 1997;(339):71–5.
26. Dellinger EP, Caplan ES, Weaver LD, et al. Duration of preventive antibiotic administration for open extremity fractures. *Arch Surg* 1988;123(3):333–9.
27. Zalavras CG, Patzakis MJ, Holtom P. Local antibiotic therapy in the treatment of open fractures and osteomyelitis. *Clin Orthop Relat Res* 2004;(427):86–93.
28. McKee MD, Wild LM, Schemitsch EH, et al. The use of an antibiotic-impregnated, osteoconductive, bioabsorbable bone substitute in the treatment of infected long bone defects: early results of a prospective trial. *J Orthop Trauma* 2002;16(9):622–7.
29. Penn-Barwell JG, Murray CK, Wenke JC. Local antibiotic delivery by a bioabsorbable gel is superior to PMMA bead depot in reducing infection in an open fracture model. *J Orthop Trauma* 2014;28(6):370–5.
30. Holtom PD, Warren CA, Greene NW, et al. Relation of surface area to in vitro elution characteristics of vancomycin-impregnated polymethylmethacrylate spacers. *Am J Orthop* 1998;27:207–10.
31. Baker AS, Greenham LW. Release of gentamicin from acrylic bone cement. Elution and diffusion studies. *J Bone Joint Surg Am* 1988;70(10):1551–7.
32. Greene N, Holtom PD, Warren CA, et al. In vitro elution of tobramycin and vancomycin polymethylmethacrylate beads and spacers from simplex and palacos. *Am J Orthop* 1998;27(3):201–5.
33. Torholm C, Lidgren L, Lindberg L, et al. Total hip joint arthroplasty with gentamicin-impregnated cement. A clinical study of gentamicin excretion kinetics. *Clin Orthop Relat Res* 1983;(181):99–106.
34. Adams K, Couch L, Cierny G, et al. In vitro and in vivo evaluation of antibiotic diffusion from antibiotic-impregnated polymethylmethacrylate beads. *Clin Orthop Relat Res* 1992;(278):244–52.
35. Ostermann PA, Seligson D, Henry SL. Local antibiotic therapy for severe open fractures. A review of 1085 consecutive cases. *J Bone Joint Surg Br* 1995;77(1):93–7.
36. Craig J, Fuchs T, Jenks M, et al. Systematic review and meta-analysis of the additional benefit of local prophylactic antibiotic therapy for infection rates in open tibia fractures treated with intramedullary nailing. *Int Orthop* 2014;38(5):1025–30.
37. Tetsworth K, Cierny G 3rd. Osteomyelitis debridement techniques. *Clin Orthop Relat Res* 1999;(360):87–96.

38. Swiontkowski MF. Criteria for bone debridement in massive lower limb trauma. *Clin Orthop Relat Res* 1989;(243):41–7.
39. Harley BJ, Beaupre LA, Jones CA, et al. The effect of time to definitive treatment on the rate of nonunion and infection in open fractures. *J Orthop Trauma* 2002; 16(7):484–90.
40. Pollak AN, Jones AL, Castillo RC, et al. The relationship between time to surgical debridement and incidence of infection after open high-energy lower extremity trauma. *J Bone Joint Surg Am* 2010;92(1):7–15.
41. Schenker ML, Yannascoli S, Baldwin KD, et al. Does timing to operative debridement affect infectious complications in open long-bone fractures? A systematic review. *J Bone Joint Surg Am* 2012;94(12):1057–64.
42. Skaggs DL, Friend L, Alman B, et al. The effect of surgical delay on acute infection following 554 open fractures in children. *J Bone Joint Surg Am* 2005;87(1): 8–12.
43. Weber D, Dulai SK, Bergman J, et al. Time to initial operative treatment following open fracture does not impact development of deep infection: a prospective cohort study of 736 subjects. *J Orthop Trauma* 2014;28(11):613–9.
44. Bhandari M, Adili A, Schemitsch EH. The efficacy of low-pressure lavage with different irrigating solutions to remove adherent bacteria from bone. *J Bone Joint Surg Am* 2001;83(3):412–9.
45. Anglen JO. Comparison of soap and antibiotic solutions for irrigation of lower-limb open fracture wounds. A prospective, randomized study. *J Bone Joint Surg Am* 2005;87(7):1415–22.
46. Bhandari M, Adili A, Lachowski RJ. High pressure pulsatile lavage of contaminated human tibiae: an in vitro study. *J Orthop Trauma* 1998;12(7):479–84.
47. Owens BD, White DW, Wenke JC. Comparison of irrigation solutions and devices in a contaminated musculoskeletal wound survival model. *J Bone Joint Surg Am* 2009;91(1):92–8.
48. Dirschl DR, Duff GP, Dahners LE, et al. High pressure pulsatile lavage irrigation of intraarticular fractures: effects on fracture healing. *J Orthop Trauma* 1998;12(7): 460–3.
49. FLOW Investigators, Bhandari M, Jeray KJ, et al. A trial of wound irrigation in the initial management of open fracture wounds. *N Engl J Med* 2015;373(27): 2629–41.
50. Weitz-Marshall AD, Bosse MJ. Timing of closure of open fractures. *J Am Acad Orthop Surg* 2002;10(6):379–84.
51. Patzakis MJ. Clostridial myonecrosis. *Instr Course Lect* 1990;39:491–3.
52. DeLong WG Jr, Born CT, Wei SY, et al. Aggressive treatment of 119 open fracture wounds. *J Trauma* 1999;46(6):1049–54.
53. Jenkinson RJ, Kiss A, Johnson S, et al. Delayed wound closure increases deep-infection rate associated with lower-grade open fractures: a propensity-matched cohort study. *J Bone Joint Surg Am* 2014;96(5):380–6.
54. Hohmann E, Tetsworth K, Radziejowski MJ, et al. Comparison of delayed and primary wound closure in the treatment of open tibial fractures. *Arch Orthop Trauma Surg* 2007; 127(2):131–6.
55. Patzakis MJ, Wilkins J, Moore TM. Considerations in reducing the infection rate in open tibial fractures. *Clin Orthop Relat Res* 1983;(178):36–41.
56. Stannard JP, Volgas DA, Stewart R, et al. Negative pressure wound therapy after severe open fractures: a prospective randomized study. *J Orthop Trauma* 2009; 23(8):552–7.

57. Gopal S, Majumder S, Batchelor AG, et al. Fix and flap: the radical orthopaedic and plastic treatment of severe open fractures of the tibia. *J Bone Joint Surg Br* 2000;82(7):959–66.
58. Pollak AN, McCarthy ML, Burgess AR. Short-term wound complications after application of flaps for coverage of traumatic soft-tissue defects about the tibia. The Lower Extremity Assessment Project (LEAP) Study Group. *J Bone Joint Surg Am* 2000;82-A(12):1681–91.
59. Cierny G 3rd, Byrd HS, Jones RE. Primary versus delayed soft tissue coverage for severe open tibial fractures. A comparison of results. *Clin Orthop Relat Res* 1983;(178):54–63.
60. Godina M. Early microsurgical reconstruction of complex trauma of the extremities. *Plast Reconstr Surg* 1986;78(3):285–92.
61. Worlock P, Slack R, Harvey L, et al. The prevention of infection in open fractures: an experimental study of the effect of fracture stability. *Injury* 1994;25(1):31–8.
62. Brumback RJ, Ellison PS Jr, Poka A, et al. Intramedullary nailing of open fractures of the femoral shaft. *J Bone Joint Surg Am* 1989;71(9):1324–31.
63. Bhandari M, Guyatt G, Tornetta P 3rd, et al. Randomized trial of reamed and unreamed intramedullary nailing of tibial shaft fractures. *J Bone Joint Surg Am* 2008; 90(12):2567–78.
64. Schemitsch EH, Kowalski MJ, Swiontkowski MF, et al. Cortical bone blood flow in reamed and unreamed locked intramedullary nailing: a fractured tibia model in sheep. *J Orthop Trauma* 1994;8(5):373–82.
65. Schemitsch EH, Kowalski MJ, Swiontkowski MF, et al. Comparison of the effect of reamed and unreamed locked intramedullary nailing on blood flow in the callus and strength of union following fracture of the sheep tibia. *J Orthop Res* 1995; 13(3):382–9.
66. Benirschke SK, Agnew SG, Mayo KA, et al. Immediate internal fixation of open, complex tibial plateau fractures: treatment by a standard protocol. *J Orthop Trauma* 1992;6(1):78–86.
67. Kregor PJ, Stannard JA, Zlowodzki M, et al. Treatment of distal femur fractures using the less invasive stabilization system: surgical experience and early clinical results in 103 fractures. *J Orthop Trauma* 2004;18(8):509–20.
68. Moed BR, Kellam JF, Foster RJ, et al. Immediate internal fixation of open fractures of the diaphysis of the forearm. *J Bone Joint Surg Am* 1986;68(7):1008–17.
69. Pape HC, Tornetta P 3rd, Tarkin I, et al. Timing of fracture fixation in multitrauma patients: the role of early total care and damage control surgery. *J Am Acad Orthop Surg* 2009;17(9):541–9.
70. Edwards CC, Simmons SC, Browner BD, et al. Severe open tibial fractures. Results treating 202 injuries with external fixation. *Clin Orthop Relat Res* 1988;(230):98–115.
71. Marsh JL, Nepola JV, Wuest TK, et al. Unilateral external fixation until healing with the dynamic axial fixator for severe open tibial fractures. *J Orthop Trauma* 1991; 5(3):341–8.
72. Sirkin M, Sanders R, DiPasquale T, et al. A staged protocol for soft tissue management in the treatment of complex pilon fractures. *J Orthop Trauma* 1999; 13(2):78–84.